

Problems with developmental stability in two rodent species from Chernobyl

T.K. Oleksyk^{1,2}, M.H. Smith^{1,2}, S.P. Gashchak³, J.M. Novak^{1,2} and J.R. Purdue⁴

¹ The University of Georgia's, Savannah River Ecology Laboratory, Drawer E, Aiken, SC 29802, U.S.A.

² Institute of Ecology, The University of Georgia, Athens, GA 30602, U.S.A.

³ International Radioecology Laboratory, Slavutych 07100, Ukraine

⁴ Illinois State Museum, 1011 East Ash St., Springfield, IL 62703, U.S.A.

Abstract. Changes in developmental patterns are some of the most important effects that may be observed at radioactively contaminated sites like those at Chernobyl. Developmental instability may arise from the interactions between an organism's genotype and its environment and be manifested as deviant morphology. Fluctuating asymmetry (FA) is a measure of deviations from the expected bilateral symmetry of the body. Our objective was to test for differences in FA in two rodent species (*Apodemus flavicollis* and *Clethrionomys glareolus*) that live in habitats surrounding the failed Chernobyl reactor. Rodents were collected from four different locations (two contaminated and two reference sites), and a series of adult skulls were photographed and 24 landmarks on each skull were located on digital images from photographed skulls. FA was calculated from the differences in these landmarks on the right and left side of the skull. Significantly more asymmetry (~2X) was observed in mice and voles living around Chernobyl compared to those living at the uncontaminated reference sites. These are relatively large effects in comparison to those previously found for plants and swallows. FA can be a cheap, easily determined and sensitive indicator of radiation-induced stress in small mammals. FA can be used to prioritize environments for remediation efforts and to efficiently evaluate the effectiveness of the remediation efforts.

1. INTRODUCTION

Fluctuating asymmetry (FA) is an estimate of small and non-directional departures from the expected bilateral symmetry [10]. FA provides an estimate of developmental stability, because the two sides of a bilaterally symmetrical organism would share the same underlying genotype, and therefore are expected to be identical in a favorable environment [6,9]. FA is expected to be higher in populations exposed to significant levels of stress caused by anthropogenic contamination [6]. FA can be estimated as the variance of a distribution among individuals of the differences between the left and right sides.

Environmental radiation has imposed a significant level of stress on populations of organisms living in the landscapes contaminated after the 1986 Chernobyl meltdown. However, mammalian populations from Chernobyl have not been tested for FA despite the fact that they live in the most contaminated areas around the failed reactor [1]. In the absence of data on humans, small mammal populations can be used as a model for evaluating risks of radioactive contamination in human populations, because they are relatively sensitive to radioactivity.

FA can only be used as a reliable indicator of environmental quality when a substantial number of reference populations with little or no known contamination are sampled. In this case, it is essential to establish the background level of FA in a species using at least two reference populations with no known contamination [7]. In addition, if FA is determined mostly by environmental factors, different species should express similar levels of FA in the same localities if they receive similar amounts of radioactive exposure.

Our overall objective is to determine whether asymmetry is elevated in radioactively contaminated populations compared to that in reference populations of two rodents species

common around the Chernobyl Nuclear Power Plant. Individual estimates for FA of size and shape of skulls were calculated for each of 3 populations of *Apodemus flavicollis* and 3 populations of *Clethrionomys glareolus*. For each of the species we used a highly contaminated population from Chernobyl, and two reference populations located north and south from the radioactive area. We then tested for differences among populations in the amounts of FA.

2. MATERIALS AND METHODS

We collected 45 individuals of *Apodemus flavicollis* and 47 individuals of *Clethrionomys glareolus* from 3 locations in Ukraine (Table 1). The first population, trapped near Gluboke Lake, came from within the highly contaminated 10 km zone in close proximity to the failed reactor. The captured animals at this site had large amounts of intramuscular contamination by ^{137}Cs [8]. One control population came from a site approximately 350 km southwest of the failed reactor. This population, Lysychynsi, had no known radioactive contamination. Another control population, Vyshenky, came from an uncontaminated location 200 km northeast of Chernobyl.

Our study concentrated on the fluctuating asymmetry (FA) of skulls. Each specimen was cleaned with dermestid beetles and dried. We took pictures of the ventral surface of each skull with a 35mm camera using a ring flash to provide even lighting. Two pictures of each skull were taken to account for the effects of placement on the measurement error. Photographs were developed and scanned. Twenty-four evolutionary homologous landmarks were chosen on each side of the skull to represent its entire surface in the same manner for both of the species. Landmarks were independently placed on each of the pictures twice in a digitizing program TPSDIG [12]. All of the statistical analyses were conducted using SAS 8.1 [14].

The FA of size and shape were estimated as described in Klingenberg and McIntyre [3]. Asymmetry in overall size was estimated using centroid size (CS). The CS for each side of each picture for each individual skull was calculated as the square root of the sum of squared distances from a set of landmarks to their centroid [15]. Each skull was scaled to the unit CS in the analysis to eliminate the effect of individual size.

We analyzed the FA of shape by superimposing the configurations of landmarks using the Procrustes method [3,13]. First, landmark configurations of the left sides of the skulls are reflected to their mirror images to align corresponding landmarks of right and left sides. After configurations are scaled to unit CS, the points with average coordinates and configurations are rotated around their centroid to achieve the best fit. The output of the Procrustes procedure contains the coordinates of superimposed landmarks. Asymmetry can then be measured as the deviations between the pairs of the corresponding superimposed landmarks.

To estimate the FA of total skull size, we used an ANOVA with CS as the dependent variable, side as a fixed effect and individuals as a random effect as recommended by Palmer [9]. To assess the FA of skull shape, we calculated the overall sums of squares for each of the main effects, interaction term and the error by adding the individual sums of squares for each of the effects across the X and Y coordinates [3]. The interaction term in this model represents the variation in left-right differences among individuals, which is a measure of FA. The measurement error was used to test for the significance of the individual*side interaction effect. The variance component of the interaction term provides an unbiased estimate of FA in each population. Degrees of freedom were calculated using the Satterthwaite approximation as suggested by Palmer [9]. Degrees of freedom for the Procrustes ANOVA were the degrees of freedom for each of the effects multiplied by the number of landmark coordinates minus four [3]. Differences in individual population FAs were tested using pairwise *F*-tests. All probabilities

generated from *F*-tests were tested with a sequential Bonferroni procedure as suggested by Palmer [9].

3. RESULTS

Normality of frequency distributions of CS for each population and each species were addressed using Kolmogorov-Smirnov tests of the frequency distribution of the centroid sizes compared to an expected normal distribution. The frequency distributions of our data for each population were inspected for the presence of bimodality (antisymmetry) or unusual outliers. After adjusting the overall error rate to the 0.05 level, there were no significant deviations from normality as indicated by the Kolmogorov-Smirnov test.

Measurement error was addressed during *F*-tests. We were testing whether our FA estimate was significantly larger than predicted due to error alone. In Vyshenky it was not possible to calculate a reliable estimate of the FA of shape compared to the relatively high measurement error for *A. flavicollis*. We also failed to calculate the significant estimate of FA of shape in *C. glareolus* from Gluboke Lake. All of the other measurements of FA of size and shape were statistically significant ($p \leq 0.05$; Table 1).

FA of size was significantly greater than the variance expected due to the measurement error for all populations except that of *A. flavicollis* in Vyshenky. Thus, this population was not used in the pairwise comparison. FA of size was significantly higher at Gluboke Lake than in the reference populations for *A. flavicollis* (Table 1). FA of shape was not different between populations of *C. glareolus* in Lysychynsi and Vyshenky. Measurements of FA of size were always higher in populations of *C. glareolus* than in *A. flavicollis* from the same locations.

FA of shape was significantly larger than the variance expected due to measurement error for all the populations. The FAs of shape in Gluboke Lake was significantly greater in FA compared to the other two populations both for *A. flavicollis* and *C. glareolus* (Table 1, Fig. 1,2). Measurements of FA of shape also did not differ from each other at the same locations between the two different species ($p \geq 0.05$, Table 1).

4. DISCUSSION

Fluctuating asymmetry (FA) refers to the difference between the right and left sides in characters that should otherwise be bilaterally symmetrical. It is likely to be a consequence of epigenetic stress that affects individuals during development. Ionizing radiation should impose stress on individual animals in the Chernobyl contaminated landscapes. We predicted that higher values of FA would be observed in animals of different species of small mammals from the contaminated site closest to the failed reactor as compared to those from the uncontaminated reference sites.

Overall, a higher level of FA was documented for the population in closest proximity to the failed Chernobyl reactor for the asymmetry of both size and shape in two species of small mammals. Fluctuating asymmetry of shape was higher in the population from Gluboke Lake than in two reference populations outside the contaminated site. In turn, reference populations were not different from each other. FA of size was always higher for *C. glareolus* than for *A. flavicollis*. Finally, levels of asymmetry of shape were the same in two different species from the same locations.

FA has been extensively used as a measure of developmental stability because of its presumed environmental basis. However, the environmental influence of stress on developmental instability and consequently on FA as its measure is not straightforward. Genetic components of FA are generally low, whereas the environmental components are usually high.

Table 1: Estimated values of the fluctuating asymmetry (FA) of size and shape for two different species of small mammals from a population in Chernobyl (Gluboke Lake) and two reference populations (Lysychyntsi and Vyshenky). Significant values at $p \leq 0.05$ are underlined. Degrees of freedom are calculated with Satterthwaite procedure. Differences between populations are tested with pairwise F -tests. P -values for the pairwise comparisons are Bonferroni corrected. Populations that were significantly different in the pairwise FA comparisons are indicated in the last column. Values that are significantly different are marked by different letters (F -test, $A \neq B$, $p \leq 0.05$).

FA of Size	<i>Clethrionomys glareolus</i>				<i>Apodemus flavicollis</i>			
	<i>N</i>	<i>d.f.</i>	FA values	Pairwise Comparisons	<i>N</i>	<i>d.f.</i>	FA values	Pairwise Comparisons
1. Lysychyntsi	15	14	<u>2.159×10^{-07}</u>	A	15	11	<u>1.07×10^{-08}</u>	A
2. Gluboke Lake	12	11	<u>2.576×10^{-07}</u>	not tested	13	7	<u>4.05×10^{-08}</u>	B
3. Vyshenky	20	19	<u>1.372×10^{-07}</u>	A	17	1	<u>1.49×10^{-08}</u>	not tested
FA of shape								
1. Lysychyntsi	15	352	<u>8.766×10^{-08}</u>	A	15	329	<u>7.49×10^{-08}</u>	A
2. Gluboke Lake	12	234	<u>1.556×10^{-07}</u>	B	13	366	<u>1.64×10^{-07}</u>	B
3. Vyshenky	20	489	<u>7.512×10^{-08}</u>	A	17	457	<u>9.36×10^{-08}</u>	A

Similar estimates of FA of shape in two rodent species are consistent with the hypothesis that FA has mostly environmental origins. On the other hand, FA of size was always higher for *C. glareolus* than *A. flavicollis*. This finding may reflect differences in the specifics of the natural history for these two species. However, what causes the difference to appear in the size asymmetry but not in the shape asymmetry is not clear. Our study demonstrated an almost two fold increase in FA of shape in the contaminated populations from Chernobyl area for two species of small mammals as compared to their reference populations. Many other field studies have linked toxic agents in contaminated areas and demonstrated FA as an indicator of environmental quality. Radiation from Chernobyl affected levels of FA in three species of plants in areas near the Chernobyl Exclusion Zone [5]. Several studies demonstrated plants increase their FAs close to sources of aerial contamination [2,4]. In mammals, high heavy metal concentrations were correlated with high FA values in common shrews [11]. Gray seals have also been reported to have higher levels of FA in highly polluted areas versus those in relatively pristine reference areas [16].

In conclusion, FA of skulls in *A. flavicollis* mice may represent an indicator of the level of radioactive contamination in the animal's environment. Highly contaminated populations of a small mammal species expressed significantly higher levels of FA calculated both as asymmetry of size and shape in two species of small mammals. Fluctuating asymmetry of shape was higher in the population from Gluboke Lake then in two reference populations south and north from the contaminated site. FA of size was always higher for *C. glareolus* than *A. flavicollis*. Finally, levels of asymmetry of shape were the same in two different species from the same locations. These are relatively large effects in comparison to those previously found for plants and swallows. FA can be a cheap, easily determined and sensitive indicator of radiation-induced stress in small mammals. FA can be used to prioritize environments for remediation efforts and to efficiently evaluate the effectiveness of remediation efforts.




Figure 1: Fluctuating asymmetry (FA) estimates in *Clethrionomys glareolus*. Values that are significantly different are marked by different letters separately for FA of size and FA of shape. Values that are significantly different are marked by different letters (F -test, $A \neq B$, $p \leq 0.05$). Value of FA of size was not significantly different from zero due to the large amounts of variance other than FA at Gluboke Lake ($p \geq 0.05$, ns).




Figure 2: Fluctuating asymmetry (FA) estimates in *Apodemus flavicollis*. Values that are significantly different are marked by different letters separately for FA of size and FA of shape. Values that are significantly different are marked by different letters (F -test, $A \neq B$, $p \leq 0.05$). Value of FA of size was not significantly different from zero at Vyshenky ($p \geq 0.05$, ns).

Acknowledgements

Support for this research came from a contract (DE-FC09-96SR18546) between the U. S. Department of Energy and The University of Georgia. Facilities and logistical support in Ukraine were provided through Dr. Bondarkov, director of the International Radioecology Laboratory, Slavutych, Ukraine. Assistance in handling and storage of rodent skulls was provided by E. McGhee at the Georgia Museum of Natural History, Athens, GA. Other help came from C. Jagoe, T. Glenn, A. Arkhipov, I. Chizhevskiy, P. E. Johns, A. J. Majeske, and I. Shchohalevich.

References

- [1] Baker R.J., Hamilton M.J., Van Den Bussche R.A., Wiggins L.E., Sugg D.W., Smith M.H., Lomakin M.D., Gaschak S.P., Rudenskaya G.A., Buntova E.G., Chesser R.K., *Journal of Mammalogy* **77** (1996) 155-170.
- [2] Graham J.H., Siegel M.I., Anistranski J., *Genetica* **89** (1993) 121-137.
- [3] Klingenberg C.P., McIntyre G.S., *Evolution* **52** (1998) 1363-1375.
- [4] Kozlov M.V., Wiseley B.J., Koricheva J.K., Haukoja E., *Journal of Applied Ecology* **33** (1996) 1489-1495.
- [5] Møller A.P., *Oikos* **81** (1998) 444-448.
- [6] Møller A.P., Swaddle J.P., *Asymmetry, Developmental Stability and Evolution*, Oxford University Press, Oxford, 1997) 291 pp.
- [7] Oleksyk T.K., Distribution and effects of Radioactive Contamination in Rodent Populations From Chernobyl. (PhD Dissertation, The Institute of Ecology, The University of Georgia, Athens, GA, 2001) 149 pp.
- [8] Oleksyk T.K., Gaschak S.P., Glenn T.C., Jagoe C.H., Peles J.D., Purdue J.R., Tsyusko O.V., Zalissky O.O., Smith M.H., *Journal of Environmental Radioactivity*, (in press) (2001).
- [9] Palmer A.R., Fluctuating asymmetry analyses: a primer. T.A. Markow (Ed.), *Developmental Instability: It's origin and Evolutionary Implications*, (Kluwer Academic Publishers, Netherlands, 1994) pp. 335-364.
- [10] Palmer A.R., *Bioscience* **46** (1996) 518-532.
- [11] Pankakoski E., Koivisto I., Hyvarinen H., *Acta Zoologica Fennica* **191** (1992) 137-144.
- [12] Rohlf F.J., TPSDIG. (Department of Ecology and Evolution, State University of New York at Stony Brook, Stony Brook, NY, 2001.
- [13] Rohlf F.J., Slice D., *Systematic Zoology* **39** (1990) 40-59.
- [14] SAS, In, SAS Institute Inc., Cary, N.C., 2000.
- [15] Slice D.E., Bookstein F.L., Marcus L.F., Rohlf F.J., *Advances in Morphometrics. A glossary for geometric morphometrics*. L.F. Marcus, M. Corti, A. Loy, G.J.P. Naylor, D. Slice (Eds.), Vol. 284, (Plenum, New York, 1996) pp. 531-551.
- [16] Zakharov V.M., Olsson M., Yablokov A.V., Esipenko A.G., Does environmental pollution affect the developmental stability of the baltic grey seal (*Halichoerus grypus*)? A.V. Yablokov, M. Olsson (Eds.), *Influence of human activities on the Baltic ecosystem. Proceedings of the Soviet-Swedish Symposium*, (Gidrometeoizdat, Leningrad, 1989) pp. 96-108.